

**AN IMPLANTABLE INTRACRANIAL PHOTO APPLICATOR FOR LONG TERM
FRACTIONATED PHOTODYNAMIC AND RADIATION THERAPY IN THE BRAIN AND
METHOD OF USING THE SAME**

5

Abstract of the Disclosure

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

An inflatable balloon is disposed into a body cavity and inflated to expand into the body cavity. A subcutaneously implanted, resealable catheter is coupled to the inflatable balloon. The resealable catheter provides repetitive access for an optical fiber disposed through a first lumen to illuminate the inner surface and for a radiation source disposed on the distal tip of a wire to provide repetitive long term, fractionated photodynamic and radiation therapy to tissues adjacent to the inner surface. A light diffusing fluid is disposed in the inflatable balloon. The catheter has a self-healing membrane coupled to and closing its proximal end. An insert is coupled to the proximal end and to the first lumen in the subcutaneous, resealable catheter. The insert is funnel shaped, but not necessarily concentric, narrowing down to where the insert is coupled to the lumen to ease in disposition of the insert into the patient and to facilitate introduction of the optical fiber therethrough without damage to the optical fiber. The insert can be disposed into or placed on top and supported only by the bony cranium of the patient and is supported by the cranium so that forces applied to the insert are prevented from being transmitted to underlying brain tissue.